



Nomenclatural entanglements associated with *Racomitrium chlorocarpum* (Grimmiaceae)

HALINA BEDNAREK-OCHYRA

Laboratory of Bryology, Institute of Botany, Polish Academy of Sciences, ul. Lubicz 46, 31-512 Kraków, Poland, e-mail: h.bednarek@botany.pl

Abstract

A nomenclatural history of one of the most neglected and misunderstood species *Racomitrium chlorocarpum* is presented. Valid publication of this species name was incorrectly ascribed to M. Fleischer and *R. chlorocarpum* of 1904 is merely a younger isonym of *R. chlorocarpum* Paris of 1897. The species name is lectotypified with a specimen from Tasmania, Australia, deposited in the Hooker herbarium (BM), and an isolectotype is present in the herbarium of W. Mitten (NY). *Racomitrium chlorocarpum* is a typical representative of the genus *Bucklandiella*, the largest segregate of the broadly interpreted genus *Racomitrium* and the relevant transfer to this genus is effected as *Bucklandiella chlorocarpa*, *comb. nov.* The diagnostic characters of the species are briefly summarised. This is a hydrophytic species, endemic to Australasia and is known from SE Australia, Tasmania and New Zealand.

Key words: Australia, author citation, Bryophyta, *Bucklandiella*, lectotypification, New Zealand, nomenclature, Tasmania, taxonomy

Introduction

The genus *Racomitrium* Bridel (1818[1819]: 78) is well represented in Australasia, both in terms of frequency of species and cover. However, the real number of species in this region is still unknown because the genus has not been carefully and critically revised to provide a sound taxonomic framework for Australasian material. In addition, the traditionally conceived genus *Racomitrium* was split into four genera (Bednarek-Ochyra *et al.*, 2001; Ochyra *et al.*, 2003), three of which are present in Australasia, including *Racomitrium* s. str. (2 species), *Niphotrichum* Bedn.-Ochyra *et Ochyra in Ochyra et al.* (2003: 137) (2 species) and *Bucklandiella* Roivainen (1972: 116). The latter is the largest and least understood segregate of *Racomitrium*. Almost all species described in the nineteenth century from Australia and New Zealand, as well as from other regions of the Southern Hemisphere, and correctly belonging within this genus, were lumped with *Bucklandiella crispula* (Hooker & Wilson 1844: 544) Bedn.-Ochyra *et Ochyra in Ochyra et al.* (2003: 144) [as *Racomitrium crispulum* (Hook.f. et Wilson) Wilson (1854: 75)] (Dixon, 1926; Clifford, 1955; Lawton, 1973). However, various authors have shown that many of these are distinct and well defined species which are only distantly related to *B. crispula* (e.g. Roivainen, 1955; Bell, 1974; Deguchi, 1984; Bednarek-Ochyra *et al.*, 1999; Bednarek-Ochyra & Ochyra, 1998, 2010, 2011, 2012a,b,c, 2013a,b; Ochyra *et al.*, 2008a,b; Blockeel *et al.*, 2009; Ellis *et al.*, 2011a, 2012a,b, 2013a,b; Larraín *et al.*, 2011; Ochyra & van Rooy, 2013).

One of species which was merged with *Bucklandiella crispula* is *Racomitrium chlorocarpum* (Dixon, 1926). During the course of ongoing revisionary studies of Australasian *Bucklandiella* it became evident that this neglected species has an exceptionally chequered taxonomic and nomenclatural history. Although this name has often been used in the Australasian bryological literature, it is not certain who actually was the first to publish it validly. Consequently, this name has not been lectotypified, nor is the correct circumscription of this species known. For that reason the authority is abandoned on the first appearance of this name in this account and, in addition, the nomenclatural issues associated with *Racomitrium chlorocarpum* are thoroughly discussed.

Nomenclatural history of *Racomitrium chlorocarpum*

It is currently accepted (Wijk *et al.*, 1967) that *Racomitrium chlorocarpum* was validly published by Fleischer (1904: 377). When describing the new Javanese species *R. microphyllum* Fleischer (1904: 377), he stated: “Diese Art steht dem *R. chlorocarpum* (Mitt.) Par. aus Neu-Seeland nahe, welches aber verlängerte, stark buchtige Blattzellen hat”. This very brief diagnosis is utterly uninformative since elongate and strongly sinuose-walled laminal cells are typical for all species of the broadly circumscribed *Racomitrium*. Yet, it constitutes a valid description of this species, even though *R. chlorocarpum* should be considered as a kind of a *nomen subnudum* in this publication. Although Fleischer (1904) ascribed this name to “(Mitt.) Par.”, the compilers of *Index muscorum* gave “Mitt. ex Fleisch.” as the authorship of this name. They accepted that Mitten (in Mueller, 1881) was the first to propose the epithet *chlorocarpum* as a *nomen nudum* in the combination with the generic name *Dryptodon* Bridel (1826: 191) and no valid diagnosis of this species was ever published (Wijk *et al.*, 1967).

The name *Racomitrium chlorocarpum* was published for the first time in *Flora Novae Zelandiae* by Hooker (1867: 426). He described *Racomitrium crispulum* and distinguished two varieties within it: [var.] α which he diagnosed by “Leaves gradually attenuated; cells of upper portion rounded” and [var.] β having “Leaves with a suboval base, obtuse and apiculate”. At the end of the line describing the latter variety Hooker appears to be including “*R. chlorocarpum*, Mitten, mss.” as a synonym. The status of this binomial is clearly confirmed in the index to this opus where, on p. 792, *Racomitrium chlorocarpum* is printed in italics and this type is used for synonyms as explained at the beginning of the index on p. 770.

In the following paragraph dealing with the distribution of these varieties, Hooker (1867) has “Var. β . Northern Island: on alpine rocks, *Colenso*” and then the note that Mitten had distinguished “the Northern Island form as a species (*R. chlorocarpum*)”. In addition, “Tasmania and Kerguelen’s Land” are indicated as parts of the range of this variety. On the other hand, var. α is shown to occur only on “Campbell’s Island, *J.D.H.*”, so it is clearly the type variety of this species since the cited specimen evidently represents the type material of *Dryptodon crispulus* Hooker & Wilson (1844: 544), collected by J.D. Hooker on Campbell Island during the Antarctic voyage of 1839–1843.

The placement of the binomial *Racomitrium chlorocarpum* after the diagnosis of var. β could seemingly suggest that the epithet *chlorocarpum* may be used as the name of this variety following Art. 24.2 of the current Code (McNeill *et al.*, 2012). However, a closer analysis of this entry clearly shows that Art. 35.2 is more applicable in this case. Yet, because the epithet *chlorocarpum* is not immediately associated with var. β , it must be rejected as a name for this variety and, accordingly, this taxon must be considered as an unnamed variety of *R. crispulum*. This interpretation is directly confirmed by the way Hooker (1867) named varieties in other parts in this publication. A closer analysis of the text shows that he described a number of varieties for various species and for some of them he used names but some remained unnamed. Unfortunately, infraspecific taxa are not indexed in this work at all. Nevertheless, there are a few cases in this work which are very instructive and confirm the correctness of the interpretation of the varietal name for *Racomitrium crispulum* var. β . For example, Hooker (1867: p. 412) distinguished *Dicranum robustum* Hooker & Wilson (1844: 542) var. β . *pungens* (Hooker & Wilson 1844: 541) Hooker (1867: 412) and at the end of this entry he placed *Dicranum pungens* Hooker & Wilson (1844: 541) as a synonym. The same scheme is also accepted for two varieties of *Sphagnum cuspidatum* Ehrhart ex Hoffmann (1796: 22) (p. 401) and one variety of *Fissidens viridulus* (Swartz ex anon. 1801: 538) Wahlenberg (1812: 334) (p. 408).

Although the name *Racomitrium chlorocarpum* Mitt. ex Hook. f. of 1867 was not validly published and is a classical *nomen nudum*, this epithet was used several times in various contemporaneous publications. Mitten (1876a, b) used the name *Grimmia (Dryptodon) chlorocarpa* Mitten (1876a: 70; 1876b: 194) for specimens collected from subantarctic Îles Kerguelen and Marion Island. Given the view expressed above it is obvious that “*Grimmia (Dryptodon) chlorocarpa* Mitt.” does not constitute “a clear (if cryptic) indication” (Art. 38.14) that Hooker’s description of *Racomitrium crispulum* var. β . applies to *Grimmia chlorocarpa* Mitt., so it must be considered not validly published. Clearly, exactly the same applies to Tasmanian *Dryptodon chlorocarpus* Mitten ex Mueller (1881: 114), *Grimmia chlorocarpa* Mitten (1882: 56) and *Racomitrium chlorocarpum* Mitten ex Bastow (1887: 60).

The next work in which *Racomitrium chlorocarpum* appeared is *Index bryologicus* (Paris, 1898), where it is accepted as a species in its own right and its name, *Rhacomitrium chlorocarpum*, is ascribed to “(Mitt.) Paris”. Three synonyms are cited for this name, including *Dryptodon chlorocarpus* Mitt. ex F. Muell. of 1881, *Grimmia chlorocarpa* Mitt. of 1882 and *Racomitrium crispulum* var. β of Hook. f. of 1867. The placement of the latter synonym may be interpreted as a direct reference to the description of this unnamed variety given by Hooker (1867) and consequently this represents the first valid publication of *Racomitrium chlorocarpum*. From the entry in Paris’s (1898) *Index bryologicus*

one can conclude that the citation of this name is simply *Racomitrium chlorocarpum* Paris (1897: 108). It is because in Art. 46.3 the second sentence makes it clear that Paris's (1898) reference to Mitten's use of the epithet *chlorocarpus* or *chlorocarpa* does not constitute ascription since it appears in a list of synonyms.

The first edition of *Index bryologicus* is a compilation of 17 articles which were originally published separately in *Actes de la Société Linnéenne de Bordeaux* from May 1894 to May 1898, each with a separate journal pagination. The compilation was published in five parts with a consecutive pagination, usually several months later after the original publication (Stafleu & Cowan, 1983). It is a generally accepted custom that numerous nomenclatural novelties proposed in this opus are ascribed to the compilation which is available as a single massive book. However, it can happen that considering only the compilation can be misleading (Lücking *et al.*, 2007). Therefore the effective date and place of publication is given here (Paris, 1897), along with the alternative bibliographic data from the compilation (Paris, 1898).

If the above argumentation regarding the validation of *Racomitrium chlorocarpum* Paris is accepted, it means that the name *R. chlorocarpum* Mitt. ex M.Fleisch. must be considered merely as a later isonym which has no nomenclatural standing (see also Ochyra *et al.*, 2006). It is worth noting that Dixon (1926) was the only author who correctly ascribed *R. chlorocarpum* to Paris. However, he placed this name without any comment in synonymy with *R. crispulum*.

Typification of *Racomitrium chlorocarpum*

According to Art. 7.7 of the current Code (McNeill *et al.*, 2012) *Racomitrium chlorocarpum* “is to be typified by an element selected from the entire context” of Hooker's (1867) description of *R. crispulum* var. β . This author cited in the protologue the specimens from three distant regions, including the North Island of New Zealand, Tasmania and Îles Kerguelen. During the course of revisionary studies of austral taxa of *Bucklandiella* some specimens annotated by W. Mitten as *Grimmia* (*Racomitrium*) *chlorocarpa* have been located on the sheet stamped “Herbarium Hookerianum” in BM and on the sheet labelled “*Racomitrium chlorocarpum* (Mitt.) Paris” in the Mitten Herbarium in NY. Many of them are eligible for selection of a lectotype of *R. chlorocarpum*.

Of the specimens cited in the protologue no material collected by Colenso from the North Island was found. In the Wilson herbarium in BM there is only a small scrap collected by Colenso in New Zealand (No. 2074), named with a question mark *Dryptodon rupestris* Hooker & Wilson (1844: 544) and cited by Wilson (1854) and Hooker (1867) under this name. Therefore this specimen cannot be considered as potential material for lectotypification.

The only material from Îles Kerguelen which was cited by Hooker (1867) in the protologue of *Racomitrium crispulum* var. β is the specimen collected by J. D. Hooker in this archipelago from May to July 1840 and cited in his *Flora antarctica* as *Racomitrium protensum* A.Braun ex Duby (1830: 573) var. 3 (Wilson & Hooker, 1847). No other moss collections were made from this isolated archipelago until 1874 when H. N. Moseley, a naturalist to the *Challenger* Expedition, collected the specimen named by Mitten (1876a) as *Grimmia chlorocarpa*. In addition, in January 1875, A. E. Eaton recorded in this archipelago during the British Transit-of-Venus Expedition the specimen named by Mitten (1876b) as *Grimmia chlorocarpa*. Nevertheless, the latter two specimens are also not eligible for lectotypification of *R. chlorocarpum*.

The specimen collected by Hooker on Îles Kerguelen was described by Müller (1889) as *Grimmia suborthotrichacea* Müller (1883: 81) var. *robustissima* Müller (1889: 30) and this variety is identical to *Bucklandiella pachydictyon* (Cardot 1908: 113) Bedn.-Ochyra et Ochyra in Ochyra *et al.* (2003: 147) (Ochyra *et al.*, 2008b). Acceptance of this material as lectotype of *R. chlorocarpum* could seriously disturb the current nomenclature because this name could have priority over *R. pachydictyon* Cardot (1908: 113), the basionym of *Bucklandiella pachydictyon*, described in 1908 (Cardot, 1908). The latter name is firmly rooted in moss taxonomy (Ochyra *et al.*, 2008a,b; Blockeel *et al.*, 2008; Müller, 2009; Larrain, 2012) and such a change would destabilise the nomenclature.

A good number of specimens which were annotated by W. Mitten as *Grimmia chlorocarpa* originate from Tasmania and these are available in the Hooker herbarium in BM and in the Mitten herbarium in NY. In fact, these are almost all specimens cited in *Flora Tasmaniae* under the name *Racomitrium crispulum* (Wilson, 1858). One of the specimens in fine fruiting condition collected by Gunn at St. Patrick's River is here selected as lectotype of *Racomitrium chlorocarpum* (Fig. 1). At the same time this species is transferred to *Bucklandiella* Roiv. as it exhibits all diagnostic features of this genus including smooth laminal cells, short lanceolate peristome teeth irregularly divided in the distal part into 2–3 branchlets, and a smooth and dextrorse seta.




FIGURE 1. The lectotype specimen of *Racomitrium chlorocarpum* Paris selected from the Hooker herbarium (BM).

***Bucklandiella chlorocarpa* (Paris) Bednarek-Ochyra & Ochyra, comb. nov.**

Racomitrium chlorocarpum Paris, *Actes Soc. Linn. Bordeaux* 51: 108. 1897 (September) and *Index Bryol.*: 1072. 1898 (February) [*Grimmia chlorocarpa* Mitt., *J. Linn. Soc. Bot.* 15: 66, 70, 194. 1876, *nom. nud.*; *Dryptodon chlorocarpus* Mitt. ex F. Muell., *Fragm. Suppl.* 11: 114. 1881, *nom. nud.*; *Grimmia chlorocarpa* Mitt., *Trans. Proc. R. Soc. Victoria* 19: 56. 1882, *nom. nud.*; *Racomitrium chlorocarpum* Mitt. ex Bastow, *Pap. Proc. Roy. Soc. Tasmania* 1886: 60. 1867 [*chlorocarpa*], *nom. nud.*]. Type citation: [New Zealand] Northern Island: on subalpine rocks, *Colenso*. (Tasmania, Kerguelen's Land). Lectotype (*selected here*): "W. 11. Rhacomitrium crispulum Hf & W. St. Patrick's River, V.D.L. 1845 Gunn. 1608. (manu Wilson) *Grimmia* (*Rhacomitrium*) *chlorocarpa*, m. (manu Mitten)" – BM-Hooker 000962320!; isolectotype: NY-Mitten! Syntypes: (1) "W.W. Rhacomitrium crispulum Hf & W. St. Patrick's River, V.D.L. 1844 (on rock) Gunn. (manu Wilson) *Grimmia* (*Rhacomitrium*) *chlorocarpa*, m. (manu Mitten)" – BM-Hooker 000962319!; (2) "V. D^s. Land. 1844 Gunn 21. Rhacomitrium crispulum Hf & W. in Fl. Tasm. *Grimmia* (*Rhacomitrium*) *chlorocarpa*, m. (manu Mitten)" – BM-Hooker 000962321!; (3) 1312. Rhacomitrium crispulum Hf & W. Fl. Tasm. p. 181 Lake Echo. V. D^s. L. on stones 2. *J.D.H.*" – BM-Hooker 000962324!; (4) 1308. V. D^s L^d. On stones near Lake Echo Rhacomitrium crispulum Hf & W. Fl. Tasm. p. 181 Lake Echo. *Grimmia* (*Rhacomitrium*) *chlorocarpa*, m. (manu Mitten)" – BM-Hooker 000962323!; (5) [The sheet labelled: The New York Botanical Garden Herbarium of William Mitten *Rhacomitrium chlorocarpum* (Mitt.) Paris: "Inundated rock Mountain River N. Huon Tasmania Mr Oldfield" – NY-Mitten!; (6) same sheet: "Tasmania Oldfield" – NY-Mitten!; (7) same sheet: "On rocks Rivulet behind Cummings H^d. Western Mts. Tasmania Mr Archer" – NY-Mitten (2 specimens)!; (8) same sheet: "Tasmania Archer" – NY-Mitten!

Concluding remarks

Bucklandiella chlorocarpa is a very distinct species which can be easily distinguished from all congeners in Australasia by a combination of both gametophyte and sporophyte characters. The leaves have a wide, 2–3(–4)-seriate basal marginal border of hyaline, pellucid cells which is usually obscured by the distinct recurvature of the leaf margins near the base. The costa is short-excurrent as a stout, massive and brown awn. The capsules are large, ellipsoid, to 2 mm long and the peristome teeth are massive, 460–580 µm long on average and 100–125 µm wide near their base, deeply purple below and strongly papillose with large warty papillae. The spores are large, 25–30(–40) µm in diameter. This is a hydrophytic species, growing typically on wet, often inundated rocks in stream beds in SE Australia, Tasmania and New Zealand. A detailed taxonomic circumscription of this species, with detailed illustrations, will be published in a separate account.

Bucklandiella is currently represented by 13 species in Australasia. Of these, 12 species are known to occur in New Zealand, seven from Tasmania and Australia, and *B. lamprocarpa* (Müller 1849: 802) Bedn.-Ochyra et Ochyra in Ochyra et al. (2003: 145) is the only species which occurs in Australasia exclusively on subantarctic Macquarie Island (Blockeel et al., 2007), although this species has a much wider range in the Subantarctic (Bednarek-Ochyra & Ochyra, 1998; Ellis et al., 2010, 2011b) as well as in South Africa (Ochyra & van Rooy, 2013) and South America and Europe (Ochyra et al., 1988; Blockeel et al., 2002). However, it is very likely that this number may still increase with progress in ongoing monographic studies on this genus. It is evidenced by the recent discoveries of some new species to New Zealand (Bednarek-Ochyra & Ochyra, 2010, 2011) and resurrection of some species from oblivion including *B. elegans* (Müller 1898: 168) Bednarek-Ochyra & Ochyra (2010: 254) (Ellis et al., 2011a) and *B. chlorocarpa* in the present account. Moreover, several species previously known from elsewhere have been added to the moss flora of New Zealand (Blockeel et al., 2008; Ellis et al., 2011a, 2014), Tasmania (Blockeel et al., 2010; Ellis et al., 2011b) and Australia (Blockeel et al., 2008). On the other hand, some species reported from Australia (Streimann & Klanzenga, 2002), such as *B. affinis* (Weber & Mohr 1807: 127) Bedn.-Ochyra et Ochyra in Ochyra et al. (2003: 144), *B. heterosticha* (Hedwig 1801: 109) Bedn.-Ochyra et Ochyra in Ochyra et al. (2003: 145), *B. obtusa* (Bridel 1801: 290) Bedn.-Ochyra et Ochyra in Ochyra et al. (2003: 146) and *B. rupestris* (Hooker & Wilson 1844: 544) Bedn.-Ochyra et Ochyra in Ochyra et al. (2003: 147) must be excluded from the bryoflora of this continent.

Acknowledgements

This work was financially supported by the Polish National Centre of Sciences through grant No. N N 303 796 940 for the author and, partly, through the statutory fund of the Institute of Botany of the Polish Academy of Sciences. I am indebted to the Curators of the herbaria at BM and NY for kindly allowing me to study their herbarium holdings of *Bucklandiella*. Special thanks are due to Dr John McNeill, Edinburgh, for his priceless suggestions regarding nomenclatural matters, to my husband Ryszard Ochyra for critical comment on the manuscript and valuable additions and to Rod Seppelt, Hobart, Tasmania for checking the English.

References

- Anonymous. (1801) [Review] Olof Swartz, Med. Doct. &c., &c., &c. Dispositio systematica Muscorum frondosorum Sueciae, adjectis descriptionibus et iconibus specierum. 12mo pp. 112. Erlangae, apud Jo. Jac. Palm 1799. *Monthly Review* 34: 535–538.
- Bastow, R.A. (1887) Tasmanian mosses. *Papers and Proceedings of the Royal Society of Tasmania* 1886: 38–102.
- Bednarek-Ochyra, H. (1995) Rodzaj *Racomitrium* (Musci, Grimmiaceae) w Polsce: taksonomia, ekologia i fitogeografia [The genus *Racomitrium* (Musci, Grimmiaceae) in Poland: taxonomy, ecology and phytogeography]. *Fragmenta Floristica et Geobotanica Series Polonica* 2: 3–307. [in Polish with extensive English summary]
- Bednarek-Ochyra, H. & Ochyra, R. (1998) *Racomitrium lamprocarpum* (Müll. Hal.) Jaeg. – an addition to the moss flora of Îles Kerguelen and the Subantarctic. *Journal of Bryology* 20: 525–528.
<http://dx.doi.org/10.1179/jbr.1998.20.2.525>
- Bednarek-Ochyra, H. & Ochyra, R. (2010) *Bucklandiella allanfifei* (Grimmiaceae), a new moss species from New Zealand, with a note on South American *B. striatipila*. *Journal of Bryology* 32: 245–255.

<http://dx.doi.org/10.1179/037366810x12814321877381>

- Bednarek-Ochyra, H. & Ochyra, R. (2011) *Bucklandiella angustissima* sp. nov. (Grimmiaceae), a new austral amphipacific species with the smallest capsules and the shortest setae in the genus. *Cryptogamie, Bryologie* 32: 13–27.
<http://dx.doi.org/10.7872/cryb.v32.iss1.2011.013>
- Bednarek-Ochyra, H., Ochyra, R. & Buck, W.R. (1999) The genus *Racomitrium* (Grimmiaceae) in Brazil, with the first report of *R. subsecundum* in South America. *Brittonia* 51: 93–105.
<http://dx.doi.org/10.2307/2666564>
- Bednarek-Ochyra, H., Lamy, D. & Ochyra, R. (2001) A note on the moss genus *Codriophorus* P. Beauv. *Cryptogamie, Bryologie* 22: 105–111.
[http://dx.doi.org/10.1016/s1290-0796\(01\)01055-0](http://dx.doi.org/10.1016/s1290-0796(01)01055-0)
- Bednarek-Ochyra, H. & Ochyra, R. (2012a) Antipodal mosses: XVII. *Bucklandiella longtonii* (Bryopsida, Grimmiaceae), a new species with caducous leaf apices from subantarctic South Georgia. *Nova Hedwigia* 94: 373–384.
<http://dx.doi.org/10.1127/0029-5035/2012/0020>
- Bednarek-Ochyra, H. & Ochyra, R. (2012b) A consideration of *Bucklandiella* (Bryophyta, Grimmiaceae) in South America, with a taxonomic re-assessment of *Racomitrium looseri*. *Nova Hedwigia* 95: 153–163.
<http://dx.doi.org/10.1127/0029-5035/2012/0045>
- Bednarek-Ochyra, H. & Ochyra, R. (2012c) The taxonomic status of *Racomitrium capense* (Bryophyta, Grimmiaceae) from South Africa. *Cryptogamie, Bryologie* 33: 97–106.
<http://dx.doi.org/10.7872/cryb.v33.iss2.2012.097>
- Bednarek-Ochyra, H. & Ochyra, R. (2013a) Taxonomic status of *Grimmia procumbens* (Bryophyta, Grimmiaceae) from Australia. *Journal of Bryology* 35: 149–152.
<http://dx.doi.org/10.1179/1743282013y.0000000047>
- Bednarek-Ochyra, H. & Ochyra, R. (2013b) Diversity of Grimmiaceae subfam. Racomitrioideae in sub-Saharan Africa, including an addition of *Bucklandiella striatipila* to the moss flora of the continent. *Cryptogamie, Bryologie* 34: 3–12.
<http://dx.doi.org/10.7872/cryb.v34.iss1.2013.3>
- Bell, B.G. (1974) A synoptic flora of South Georgian mosses: V. *Willia* and *Racomitrium*. *British Antarctic Survey Bulletin* 38: 73–101.
- Blockeel, T.L., Duckett, J.G., Ordonez M.C.F., Hébrard, J.-P., Matcham, H.W., Porley, R.D., Ochyra, R., Bednarek-Ochyra, H., Soldán, Z. & Townsend, C.C. (2002) New national and regional bryophyte records, 5. *Journal of Bryology* 24: 88–91.
<http://dx.doi.org/10.1179/037366802125000403>
- Blockeel, T.L., Bednarek-Ochyra, H., Ochyra, R., Düzenli, A., Erdağ, A., Erzberger, P., Ezer, T., Hespanhol, H., Kara, R., Matteri, C.M., Müller, F., Séneca, A., Sérgio, C. & Váña, J. (2007) New national and regional bryophyte records, 15. *Journal of Bryology* 29: 139–142.
<http://dx.doi.org/10.1179/174328207x169082>
- Blockeel, T.L., Bednarek-Ochyra, H., Ochyra, R., Duckett, J.G., Erzberger, P., Hedenäs, L., Hugonnot, V., Maier, E., Marková, I., Matcham, H.W., Plášek, V., Pócs, T., Seppelt, R.D., Szűcs, P., Thouvenot, L. & van Zanten, B.O. (2008) New national and regional bryophyte records, 18. *Journal of Bryology* 30: 161–167.
<http://dx.doi.org/10.1179/174328208x282463>
- Blockeel, T.L., Bastos, C.J.P., Bednarek-Ochyra, H., Ochyra, R., Dulin, M.V., Fovet, I., Garcia, C., Hedenäs, L., Hugonnot, V., Kirmaci, M., Koponen, T., Lebouvier, M., Martins, A., Müller, F., Sabovljević, M., Lakušić, D., Schäfer-Verwimp, A., Sérgio, C., Surina, B. & Yayintaş, Ö.T. (2009) New national and regional bryophyte records, 22. *Journal of Bryology* 31: 201–210.
<http://dx.doi.org/10.1179/037366809x12469790518367>
- Blockeel, T.L., Bednarek-Ochyra, H., Cykowska, B., Ochyra, R., Düzenli, A., Ezer, T., Holyoak, D.T., Hugonnot, V., Kara, R., Larrain, J., Lebouvier, M., Preston, C.D., Schäfer-Verwimp, A., Smith, V.R., Spitale, D., Ștefănuț, S. & Váña, J. 2010. New national and regional bryophyte records, 23. *Journal of Bryology* 32: 140–147.
<http://dx.doi.org/10.1179/174328208x300688>
- Bridel, S.E. (1801) Animadversiones in Muscologiae recentiorum Tomum secundum, ab ipso auctore propositae. *Journal für die Botanik* 1800(2): 268–299.
- Bridel, S.E. (1818) *Methodus nova muscorum ad naturae normam melius instituta et Muscologiae recentiorum accommodata*. Gothae, apud A. Ukertum, xviii + 220 [+ 221, errata].
<http://dx.doi.org/10.5962/bhl.title.81816>
- Bridel, S.E. (1826–1827) *Bryologia univversa seu systematica ad novam methodum dispositio, historia et descriptio omnium muscorum frondosorum hucusque cognitorum cum synonymia ex auctoribus probatissimis*. Volumen primum. Lipsiae, sumtibus Joan. Ambros. Barth, 856 [+ 857–860] pp.
<http://dx.doi.org/10.5962/bhl.title.60260>

- Cardot, J. (1908) *La flore bryologique des terres magellaniques, de la Géorgie du Sud et de l'Antarctide*. In: *Wissenschaftliche Ergebnisse der Schwedischen Südpolar-Expedition 1901–1903 unter Leitung von Dr. Otto Nordenskjöld*. 4(8). Stockholm; Lithographisches Institut des Generalstabs, pp. 1–298 + pls. 1–11.
- Clifford, H.T. (1955) On the distribution of *Racomitrium crispulum* (H.f. & W.) H.f. & W. *The Bryologist* 58: 330–334.
[http://dx.doi.org/10.1639/0007-2745\(1955\)58\[330:otdorc\]2.0.co;2](http://dx.doi.org/10.1639/0007-2745(1955)58[330:otdorc]2.0.co;2)
- Deguchi, H. (1984) Studies on some Patagonian species of Grimmiaceae (Musci, Bryophyta). In: Inoue, H. (Ed.) *Studies on cryptogams in southern Chile*. Tokyo, Kenseisha, Tokyo, pp. 17–72.
- Dixon, H.N. (1926) Studies in the bryology of New Zealand, with special reference to the herbarium of Robert Brown, of Christchurch, New Zealand. Part IV. *New Zealand Institute Bulletin* 3: 153–238 + pl. ix.
<http://dx.doi.org/10.5962/bhl.title.57893>
- Duby, J.E. (1830) *Aug. Pyrami de Candolle Botanicon gallicum seu synopsis plantarum in flora gallica descriptarum*. Ed. 2. 2. Paris, V° Desray, pp. i–vi + 545–1068 + i–lviii.
<http://dx.doi.org/10.5962/bhl.title.6352>
- Ellis, L.T., Asthana, A.K., Sahu, V., Bednarek-Ochyra, B.H., Ochyra, R., Cano, M.J., Costa, D.P., Cykowska, B., Ochyra, R., Philippov, D.A., Dulin, M.V., Erzberger, P., Lebouvier, M., Mohamed, H., Ochyra, R., Orgaz, J.D., Phephu, N., van Rooy, J., Stebel, A., Suárez, G.M., Schiavone, M.M., Townsend, C.C., Váňa, J., Vončina, G., Yayintaş, Ö.T., Yong, K.T. & Zander, R.H. (2010) New national and regional bryophyte records, 25. *Journal of Bryology* 32(4): 311–322.
- Ellis, L.T., Bednarek-Ochyra, H., Ochyra, R., Calvo Aranda, S., Colotti, M.T., Schiavone, M.M., Dulin, M.V., Erzberger, P., Ezer, T., Kara, R., Gabriel, R., Hedenäs, L., Holyoak, D.T., Ódor, P., Papp, B., Sabovljević, M., Seppelt, R.D., Smith, V.R., Sotiaux, A., Szurdoki, E., Vanderpoorten, A., van Rooy, J. & Żarnowiec, J. (2011a) New national and regional bryophyte records, 26. *Journal of Bryology* 33: 66–73.
<http://dx.doi.org/10.1179/1743282010y.0000000014>
- Ellis, L.T., Asthana, A.K., Sahu, V., Srivastava, A., Bednarek-Ochyra, H., Ochyra, R., Chlachula, J., Colotti, M.T., Schiavone, M.M., Hradilek, Z., Jimenez, M.S., Klama, H., Lebouvier, M., Natcheva, R., Pócs, T., Porley, R.D., Sérgio, C., Sim-Sim, M., Smith, V.R., Söderström, L., Ștefănuț, S., Suárez, G.M. & Váňa, J. (2011b) New national and regional bryophyte records, 28. *Journal of Bryology* 33: 237–247.
<http://dx.doi.org/10.1179/1743282011y.0000000022>
- Ellis, L.T., Bednarek-Ochyra, H., Cykowska, B., Ochyra, R., Garcia, C., Sérgio, C., Lebouvier, M., Manolaki, P., Giannouris, E., Kadis, C., Marková, I., Papp, B., Szurdoki, E., Peralta, D.F., Plášek, V., Ristow, R., Sabovljević, M., Sim-Sim, M., Smith, V.R., Tsakiri, E., Váňa, J., Virchenko, V.M. & Barsukov, O.O. (2012a) New national and regional bryophyte records, 30. *Journal of Bryology* 34: 45–51.
<http://dx.doi.org/10.1179/1743282011y.0000000042>
- Ellis, L.T., Alegro, A., Bednarek-Ochyra, H., Ochyra, R., Bergamini, A., Cogoni, A., Erzberger, P., Górski, P., Gremmen, N., Hespanhol, H., Vieira, C., Kurbatova, L.E., Lebouvier, M., Martinčić, A., Asthana, A.K., Gupta, R., Nath, V., Natcheva, R., Ganeva, A., Özdemir, T., Batan, N., Plášek, V., Porley, R.D., Randić, M., Sawicki, J., Schroder, W., Sérgio, C., Smith, V.R., Sollman, P., Ștefănuț, S., Stevenson, C.R., Suárez, G.M., Surina, B., Uyar, G. & Surina, Z.M. (2012b) New national and regional bryophyte records, 31. *Journal of Bryology* 34: 123–134.
<http://dx.doi.org/10.1179/1743282012y.0000000009>
- Ellis, L.T., Bednarek-Ochyra, H., Ochyra, R., Benjumea, M.J., Saís, L.V., Caparrós, R., Lara, F., Mazimpaka, V., Dulin, M.V., Garilleti, R., Gremmen, N., Grundling, P.-L., Heras, P., Infante, M., Huttunen, S., Ignatov, M.S., Korvenpää, T., Lebouvier, M., Lewis Smith, R.I., Lin, S.-H., Yang, J.-D., Linström, A., Plášek, V., Rosselló, J.A., Sawicki, J., van Rooy, J., Smith V.R. (2013a) New national and regional bryophyte records, 35. *Journal of Bryology* 35: 129–139.
<http://dx.doi.org/10.1179/1743282013y.0000000049>
- Ellis, L.T., Aranda, S.C., Asthana, A.K., Bansal, P., Nath, V., Sahu, V., Bayliss, J., Asthana, G., Srivastava, S., Yadav, S., Brugués, M., Cano, M.J., Dulin, M.V., Fudali, E., Fuertes, E., Gabriel, R., Pereira, F., Silva J.A.F., Gradstein, S.R., Hájková, P., Hájek, M., Heras, P., Infante, M., Lebouvier, M., Marka, J., Newsham, K.K., Ochyra, R., Pantović, J., Sabovljević, M.S., Phephu, N., van Rooy, J., Philippov D.A., Porley, R.D., Puche, F., Schäfer-Verwimp, A., Segarra-Moragues, J.G., Sérgio, C., Smith V.R., Ștefănuț, S., Váňa, J., Wigginton, M.J. (2013b) New national and regional bryophyte records, 37. *Journal of Bryology* 35: 290–305.
<http://dx.doi.org/10.1179/1743282013y.0000000073>
- Ellis, L.T., Aleffi, M., Asthana, A.K., Srivastava, A., Bakalin, V.A., Batan, N., Özdemir, T., Bednarek-Ochyra, H., Borovichev, E.A., Brugués, M., Cano, M.J., Choi, S.S., De Beer, D., Eckstein, J., Erzberger, P., Fedosov, V.E., Ganeva, A., Natcheva, R., Garcia C.A., Sérgio, C., Garilleti, R., Albertos, B., Puche, F., Gücel, S., Higuchi, M., Hugonnot, V., Hylander, K., Kirmaci, M., Aslan, G., Koponen, T., Lara, F., Mazimpaka, V., van Melick, H., Müller, F., Özenoglu Kiremit, H., Papp, B., Szurdoki, E., Plášek, V., Čihal, L., van der Pluijm, A., Poponessi, S., Mariotti, M.G., Reyniers, J., Sabovljević, M.S., Sawicki, J., Smith, V.R., Stebel, A., Ștefănuț, S., Sun, B.-

- Y., Váña, J. & Vednanzoni, R. (2014) New national and regional bryophyte records, 40. *Journal of Bryology* 36: 223–244.
<http://dx.doi.org/10.1179/1743282014y.0000000115>
- Fleischer, M. (1904) *Die Musci der Flora von Buitenzorg (zugleich Laubmoosflora von Java). Enthaltend alle aus Java bekannt gewordenen Sphagnales und Bryales, nebst kritischen Bemerkungen vieler Archipelarten, sowie indischer und australischer Arten.* 1. *Sphagnales; Bryales (Arthrodontei[Haplolepidae])*. Leiden, Buchhandlung and Druckerei E.J. Brill, pp. xxxi + 379 + [6] pp.
<http://dx.doi.org/10.5962/bhl.title.44870>
- Hedwig, J. (1801) *Species muscorum frondosorum descriptae et tabulis aeneis lxxvii coloratis illustratae*. Lipsiae, sumtu Joannis Ambrosii Barthii; Parisiis, apud Amand Koenig, vi + 353 + lxxii pls.
<http://dx.doi.org/10.5962/bhl.title.26>
- Hoffmann, G.F. (1796) *Deutschlands Flora oder Botanisches Taschenbuch. Zweyter Theil für das Jahr 1795. Cryptogamie*. Erlangae, Johan Jacob Palm, lxxviii + 200 [+ 39 unnumbered] pp.
- Hooker, J.D. (1867) *Handbook of the New Zealand flora: a systematic description of the native plants of New Zealand and the Chatham, Kermadec's, Lord Auckland's Campbell's and Macquarrie's Islands, Part 2*. London, Reeve & Co., pp. xli–xlvi + 393–798.
- Hooker, J.D. & Wilson, W. (1844) Musci antarctici; being characters with brief descriptions of the new species of mosses discovered during the voyage of H. M. discovery ships, Erebus and Terror, in the southern circumpolar regions, together with those of Tasmania and New Zealand. *London Journal of Botany* 3: 533–556.
- Larraín, J.B.B. (2012) Filogenia del género *Racomitrium* Brid. (Bryophyta, Grimmiaceae) y taxonomía de las especies latinoamericanas. Concepción, La tesis doctoral en botánica, Universidad de Concepción, Dirección de Postgrado, Facultad de Ciencias Naturales Oceanográficas – Programa de Doctorado en Ciencias Biológicas área Botánica, 293 pp.
- Larraín, J., Quandt, D. & Muñoz, J. (2011) *Bucklandiella araucana* (Grimmiaceae), a new species from Chile. *The Bryologist* 114: 732–743.
<http://dx.doi.org/10.1639/0007-2745-114.4.732>
- Lawton, E. (1973) *Racomitrium crispulum* and some related species. *Bulletin of the Torrey Botanical Club* 100: 230–235.
<http://dx.doi.org/10.2307/2484802>
- Lücking, R., Kalb, K., Steiger, B. & McNeill, J. (2007) (1792) Proposal to conserve the name *Phaeographis*, with conserved type, against *Creographa*, *Ectographis*, *Fleographa*, *Hymenodecton*, *Platygramma*, and *Pyrographa* (Ascomycota: Pstrographales: Graphidaceae), along with notes on the names *Graphina* and *Phaeographina*. *Taxon* 56: 1296–1299.
<http://dx.doi.org/10.2307/25065924>
- McNeill, J., Barrie, F.R., Buck, W.R., Demoulin, V., Greuter, W., Hawksworth, D.L., Herendeen, P.S., Knapp, S., Marhold, K., Prado, J., Prud'homme van Reine, W.F., Smith, G.F., Wiersema, J.H. & Turland, N.J. (2012) *International Code of Nomenclature for algae, fungi, and plants (Melbourne Code) adopted by the Eighteenth International Botanical Congress Melbourne, Australia, July 2011*. Königstein, Koeltz Scientific Books, xxx + 208 pp.
- Mitten, W. (1876a) The Musci and Hepaticae collected by H. N. Moseley, M. A., naturalist to H. M. S. 'Challenger'. *Journal of the Linnean Society Botany* 15: 59–73.
<http://dx.doi.org/10.1111/j.1095-8339.1876.tb00223.x>
- Mitten, W. (1876b) A list of the Musci and Hepaticae collected in Kerguelen's Island by Rev. A. E. Eaton, A.M. *Journal of the Linnean Society Botany* 15: 193–197.
<http://dx.doi.org/10.1111/j.1095-8339.1876.tb00238.x>
- Mitten, W. (1882) Australian mosses. *Transactions and Proceedings of the Royal Society of Victoria* 19: 49–96.
- Mueller, F. von (1881) Addimenta pro supplemento voluminis undecimi. Musci. In: *Fragmenta phytographiae Australiae*. Volume 11 Supplementum 3. Melbourne, ex Officina Joannis Ferres, pp. 107–115.
- Müller, C. (1849) *Synopsis muscorum frondosorum omnium hucusque cognitorum. Pars prima. Musci vegetationis acrocarpicae*. Berolini, sumptibus Alb. Foerster, viii + 812 pp.
<http://dx.doi.org/10.5962/bhl.title.31>
- Müller, C. (1883) Die auf der Expedition S. M. S. »Gazelle« von Dr. Naumann gesammelten Laubmoose. *Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie* 5: 76–88.
- Müller, C. (1898) Symbolae ad bryologiam Australiae. II. *Hedwigia* 37: 76–171.
- Müller, F. (2009) An updated checklist of the mosses of Chile. *Archive for Bryology* 58: 1–124.
- Müller, K. (1889) *Laubmoose (Musci frondosi)*. In: Engler, A. (Ed.) *Die Forschungsreise S.M.S. „Gazelle“ in den Jahren 1874 bis 1876 unter Kommando des Kapitäns zur See Freiherrn von Schleinitz*. 4. Botanik. Berlin, Ernst Siegfried Mittler und Sohn, pp. 1–64.
- Ochyra, R. & van Rooy, J. (2013) Distribution of *Bucklandiella lamprocarpa* (Grimmiaceae, Musci) in South Africa. *Cryptogamie, Bryologie* 34: 359–366.
<http://dx.doi.org/10.7872/cryb.v34.iss3.2013.359>
- Ochyra, R., Sérgio, C. & Schumacker, R. (1988) *Racomitrium lamprocarpum* (C. Muell.) Jaeg., an austral moss disjunct in Portugal, with

- taxonomic and phytogeographic notes. *Bulletin du Jardin Botanique National de Belgique* 58: 225–258.
<http://dx.doi.org/10.2307/3668409>
- Ochyra, R., Żarnowiec, J. & Bednarek-Ochyra, H. (2003) *Census catalogue of Polish mosses*. Kraków, Polish Academy of Sciences, Institute of Botany, 372 pp.
- Ochyra, R., Isoviita, P., Żarnowiec, J. & Bednarek-Ochyra, H. (2006) Correct author citations for class and some subclass names of the Bryophyta. *Annales Botanici Fennici* 43: 118–122.
- Ochyra R., Bednarek-Ochyra, H. & Lewis Smith, R.I. (2008a) New and rare moss species from the Antarctic. *Nova Hedwigia* 87: 457–477.
<http://dx.doi.org/10.1127/0029-5035/2008/0087-0457>
- Ochyra, R., Lewis Smith, R.I. & Bednarek-Ochyra, H. (2008b) *The illustrated moss flora of Antarctica*. Cambridge, Cambridge University Press, xvii + 685 pp. [+ 24 pp. with plates].
- Paris, É.G. (1897) *Index bryologicus sive enumeratio muscorum hucusque cognitorum adjunctis synonyma distributioneque geographica locupletissimis. Part 1. PLA–RHY*. [Série 6] *Actes de la Société Linnéenne de Bordeaux* 51(1): 1–160.
<http://dx.doi.org/10.5962/bhl.title.643>
- Paris, É.G. (1898) *Index bryologicus sive enumeratio muscorum hucusque cognitorum adjunctis synonyma distributioneque geographica locupletissimis. 4. Parisiis, Paul Klinksieck*, pp. 965–1284.
<http://dx.doi.org/10.5962/bhl.title.643>
- Roivainen, H. (1955) Contribution to the Fuegian species of the genus *Racomitrium* Bridel. *Archivum Societatis Zoologicae Botanicae Fennicae “Vanamo”* 9(2): 85–98.
- Roivainen, H. (1972) *Bucklandiella* Roiv., nomen novum (Musci). *Annales Botanici Fennici* 9: 116.
- Stafleu, F.A. & Cowan, R.S. (1981) *Taxonomic literature. A selective guide to botanical publications and collections with dates, commentaries and types. Volume IV: P–Sak*. Ed. 2. Utrecht, Bohn, Scheltema & Holkema and The Hague, dr. W. Junk b.v., Publishers, ix + 1214 pp.
- Streimann, H. & Klanzenga, N. (2002) Catalogue of Australian mosses. *Flora of Australia Supplementary Series* 17: [1–5] + 1–259.
- Wahlenberg, G. (1812) *Flora lapponica exhibens plantas geographice et botanice consideratas, in Lapponia suecicis scilicet Umensi, Pitensi, Luliensi, Tornensi et Kemensi nec non Lapponis norvegicis scilicet Nordlandia et Finmarkia utraque indigenas, et itineribus annorum 1800, 1802, 1807 et 1810 denuo investigatas*. Berolini, in Taberna Libraria Scholae realis, lxvi + 550 pp. + xxx pls.
- Weber, F. & Mohr, D.M.H. (1807) *Botanisches Taschenbuch auf das Jahr 1807. Deutschland’s Kryptogamische Gewächse. Erste Abtheilung. Filices, Musci frondosi et Hepatici*. Kiel, in der akademischen Buchhandlung, xlvi + 509 pp. + xii pls.
- Wijk, R. van der, Margadant, W.D. & Florschütz, P.A. (1967) *Index muscorum*. Vol. 4 (P–S). Utrecht, Kemink en Zoon N.V., 604 pp.
- Wilson, W. (1854) Musci. In: Hooker, J.D. (Ed.) *The botany of the Antarctic voyage of H. M. Discovery ships Erebus and Terror in the years 1839–43, under the command of Captain Sir James Clark Ross, Kt., R.N., F.R.S. Vol. 2. Flora Novae-Zelandiae. Part. II. Flowerless plants*. London, Lovell Reeve, pp. 57–125 + pls. lxxxiii–xcii.
<http://dx.doi.org/10.5962/bhl.title.16029>
- Wilson, W. (1858) Musci. In: Hooker, J.D. (Ed.) *The botany of the Antarctic voyage of H. M. Discovery ships Erebus and Terror in the years 1839–43, under the command of Captain Sir James Clark Ross, Kt., R.N., F.R.S. Part 3. Flora Tasmaniae. Volume 2. Monocotyledones and Acotyledones*. London, Lovell Reeve, pp. [160]161–221 + pls. clxxi–clxxvii.
<http://dx.doi.org/10.5962/bhl.title.16029>
- Wilson, W. & Hooker, J.D. (1847) Musci. In: Hooker, J.D. (Ed.) *The botany of the Antarctic voyage of H. M. Discovery ships Erebus and Terror in the years 1839–43, under the command of Captain Sir James Clark Ross, Kt., R.N., F.R.S. Volume 1. Flora Antarctica. Part. II., Botany of Fuegia, the Falklands, Kerguelen’s Land, etc.* London, Reeve Brothers, pp. 395–423, 550–551 + pls. cli–clv.
<http://dx.doi.org/10.5962/bhl.title.16029>